QUEENSLAND CHILDHOOD IMMUNISATION VISUAL TOOL

H-B-VaxIIpaediatric or Engerix-B paediatric

Newborns should receive the birth dose as soon as they are medically stable, and preferably within 24 hours of birth, but the vaccine can be given within the first 7 days of life.



Beyfortus (nirsevimab)

Only recommended for infants from birth to less than 8 months of age if they meet the eligibility criteria. Nirsevimab dose is weight and age dependent.



Rotavirus

Pneumococcal* (10)

Respiratory Syncytial Virus



Prevenar 20

BCG Children <5 years living in Aboriginal and Torres Strait Islander Communities.

Bexsero

Meningococcal B*



Tuberculosis*

Infanrix hexa OR Vaxelis PRESCRIPTION ONLY MEDICINE Infanrix

Diphtheria, Tetanus, Pertussis (Whooping Cough),

Hepatitis B, Polio, Haemophilus Influenzae Type B (HIB)

PRESCRIPTION ONLY MEDICINE 🔘 Vaxelis. sanofi MCM

Rotarix The first dose must be given by 14 weeks and 6 days of age. The **second dose** must be given by 24 weeks and 6 days. Do not administer dose 2 if dose 1 was missed

Rotartx Grall Expension Suppersion Suppersion Suppersion Suppersion oral

Pneumococcal

Meningococcal B

MONTHS

9

Infanrix hexa OR Vaxelis PRESCRIPTION ONLY MEDICINE

🔾 Vaxelis.

Diphtheria, Tetanus, Pertussis (Whooping Cough),

Hepatitis B, Polio, Haemophilus Influenzae Type B (HIB)

PRESCRIPTION ONLY MEDICINE sanofi MCM Single-dose 0.5 mL pre-filled syringe without r

Influenza

Prevenar 20 Aboriginal and Torres Strait Islander children and/ or children with specified medical risk conditions only.

Bexsero Children with specified medical risk conditions.

BEXSERO Multicomponent Meningococcal group B Vaccine (recombinant, adsorbed) Suspension for injection Each pre-filled syringe contains 0.5 mL of suspension for injection (1 dose) 1 pre-filled syringe without needle. Deep intramuscular injection only.

Quadrivalent Influenza Virus (QIV) vaccine 2025 formulations: Vaxigrip Tetra

FluQuadri

Nimenrix

Infanrix



M-M-R II OR Priorix

PRESCRIPTION ONLY MEDICINE KEEP OUT OF REACH OF CHILDREN **☑ FluQuadri**"

Measles, Mumps, Rubella

Administer annually. For children aged 6 months to less than 9 years of age in the first year of administration, give 2 doses at a minimum of 4 weeks apart. Give one dose annually in subsequent years.

Bexsero

8 MONTHS TO <24 MONTHS

Beyfortus (nirsevimab)

Infants or young children with a condition associated with increased risk of severe RSV disease are eligible for an additional dose from 8 months to less than 24 months of age, ahead of their second RSV season.

The dose of nirsevimab for older children entering their second RSV season is 200mg, given as 2 intramuscular injections.

Diphtheria, Tetanus, Pertussis (Whooping Cough)



12 MONTHS



Meningococcal ACWY







Meningococcal B



Pneumococcal

Prevenar 20

H-B-Vax II paediatric or Engerix-B paediatric Low birth weight (<2000g) OR pre-term babies <32 weeks

Respiratory Syncytial Virus



Hepatitis A*

18 MONTHS

Act-HIB Act-HIB PRESCRIPTION ONLY MEDICINE KEEP OUT OF REACH OF CHILDREN Haemophilus type b Conjugate Vaccine (conjugated to tetanus protein 0 microgram) diluent for solution ± 5 mL of diluent in Pre-filled syringe with 2 separate needles Intramuscular or deep subcutaneous injection

Haemophilus Influenzae Type B (HIB)

Measles, Mumps, Rubella, Varicella Priorix-Tetra

PRESCRIPTION ONLY MEDICINE gsk GlaxoSmithKline Priorix-Tetra!

Infanrix OR Tripacel



Pneumococcal*

PRESCRIPTION ONLY MEDICINE KEEP OUT OF REACH OF CHILD TRIPACEL® 5 x 0.5 mL Single Dose Vi

Vaqta paediatric Aboriginal and Torres Strait Islander children.

Hepatitis A vaccine, Inactivated hepatitis A virus, inactivated 25 U/0.5 mL SUSPENSION FOR INTRAMUSCULAR INJECTION VADTA®
Hepatitis A vaccir
Inactivated

Diphtheria, Tetanus, Pertussis (Whooping Cough), Polio **Infanrix IPV** OR Quadracel

PRESCRIPTION ONLY MEDICINE gsk GlaxoSmithKline Infanrix

PRESCRIPTION ONLY MEDICINE KEEP OUT OF REACH OF CHILDREN Quadracel® 10 single-dose prefilled syringes 0.5 mL

Influenza*

Prevenar 20 Single dose, ONLY in children who have not previously received a dose of Prevenar 20. Refer to the Australian <u>Immunisation</u> <u>Handbook</u>

Prescription only medicin keep out of reach of child Prevenar 20 €Pfi.

Vaqta paediatric Aboriginal and Torres Strait

Islander

children.

Hepatitis A*

VAQTA inactivated 25 U/0.5 mL MSD Gegi

Quadrivalent Influenza Virus (QIV) vaccine 2025

Vaxigrip Tetra and Flucelvax Quad

formulations:

sanofi

PRESCRIPTION ONLY MEDICINE FLUCELVAX S NEEDLE-FREE AUST R 319093 CSL Seqirus

Administer annually. For children aged 6 months to less than 9 years of age in the first year of administration, give 2 doses at a minimum of 4 weeks apart. Give one dose annually in subsequent years.



25



Aboriginal and Torres Strait Islander



Medically at risk



