




























Queensland Health Immunisation Program

From 1 September 2025

NIP Funded Pneumococcal Vaccines Product Selection Guide

This product selection guide is to be used in conjunction with the Immunisation Schedule Queensland and the Australian Immunisation Handbook.

<div> <div>Australian Immunisation Handbook - Pneumococcal chapter</div> <div>Queensland Health - Immunisation Schedule Queensland</div> <div>Childhood and adolescent pneumococcal provider advice</div> </div>			<div>Prevenar 20</div>  <div>20-valent pneumococcal conjugate vaccine</div>	<div>Prevenar 13</div>  <div>13-valent pneumococcal conjugate vaccine</div>	<div>Pneumovax 23</div>  <div>23-valent pneumococcal polysaccharide vaccine</div>
<div>Infants ≤12 months</div> 	<div>All infants**</div>		 <div>Dose 1 at age 2 months (can be from 6 weeks) Dose 2 at age 4 months Dose 3 at age 12 months</div>	 <div>Not funded for use</div>	 <div>Not registered for use</div>
	<div>Aboriginal and Torres Strait Islander infants AND/OR infants with a medical risk condition**</div>		 <div>Additional dose at age 6 months</div>	 <div>Not funded for use</div>	 <div>Not registered for use</div>
<div>Children and adolescents <18 years</div> 	<div>Children and adolescents aged >12 months to <18 years with a newly diagnosed medical risk condition**</div>		 <div>Single dose at diagnosis</div>	 <div>Not funded for use</div>	 <div>Not funded for use</div>
	<div>Aboriginal and Torres Strait Islander people AND/OR people with a medical risk condition at 4 years of age OR 5 years after first Pneumovax 23 dose**</div>		 <div>Single dose ONLY required if child has not previously received a dose of Prevenar 20</div>	 <div>Not funded for use</div>	 <div>Not funded for use</div>
<div>Adults ≥18 years</div> 	<div>Adults ≥18 years with a newly diagnosed medical risk condition†</div>		 <div>Not funded for use</div>	 <div>Single dose at diagnosis</div>	 <div>Dose 1 - 1 year after Prevenar 13 Dose 2 - ≥5 years after Dose 1</div>
	<div>Aboriginal and Torres Strait Islander people ≥50 years</div>		 <div>Not funded for use</div>	 <div>Single dose</div>	 <div>Dose 1 - 1 year after Prevenar 13 Dose 2 - ≥5 years after Dose 1</div>
	<div>All adults ≥70 years</div>		 <div>Not funded for use</div>	 <div>Single dose</div>	 <div>Not funded for use</div>

*Prevenar 20 is the only pneumococcal vaccine funded for individuals under the age of 18 years.

†Refer to the Australian Immunisation Handbook for more information.

PneumococcalProductSelectionGuide_V1.1_20250811